

Comment on:
**“The Supersymmetric Ward-Takahashi Identity in 1-Loop
Lattice Perturbation Theory. I. General Procedure”**
by A. Feo; hep-lat/0305020

T. Galla^a, G. Münster^b and A. Vladikas^c

(a) Department of Physics, University of Oxford, Theoretical Physics,
1 Keble Road, Oxford OX1 3NP, UK

(b) Institut für Theoretische Physik, Universität Münster,
Wilhelm-Klemm-Str. 9, D-48149 Münster, Germany

(c) INFN, Sezione di Roma 2, c/o Dipartimento di Fisica,
Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy

Abstract

We comment on several issues concerning both the physics and authorship of article hep-lat/0305020.

In a recent paper A. Feo studies the supersymmetric Ward identities for N=1 supersymmetric Yang-Mills theory on a lattice, in the framework of lattice perturbation theory. Her article, being the first in a series of two, deals with the general procedure and the methodological foundations of the calculation. We have some critical comments on the contents of this paper.

1. The lattice SUSY Ward identity is considered in terms of the bare lattice operators, see eqs. (3.2), (3.8). From these equations eq. (3.11) does not follow, contrary to what is claimed. A multiplicative renormalization of contact terms, Fadeev-Popov term and gauge fixing term with renormalisation constants Z_{CT} , Z_{GF} and Z_{FP} could result from some of the operators B_j appearing in (3.8). In (3.11) the renormalization constants as well as the sum over all operators B_j appear. Therefore these contributions are double-counted.

2. After eq. (3.13), it is claimed that Z_S , Z_T and Z_χ are power subtraction coefficients of the operator X_S/a and therefore, based on ref. [36], their independence from a renormalisation scale is derived as a corollary. It is certainly wrong to claim that Z_χ is a power subtraction coefficient (this is true of $\tilde{m}$ in eq. (3.6)). The operator $\chi(x)$ is multiplicatively renormalisable. Thus, Z_χ has a logarithmic divergence $\ln(a\mu)$. Only in this way can the RG invariance of the product $(m_0 - \tilde{m})\chi$ (implicit in the Ward identities of the paper) be true. Since on the other hand, Z_S and Z_T are shown to be finite after eq. (3.13),

we do not understand the following statement in the conclusions: “We observe that, at least at 1-loop order in perturbation theory, Z_T is finite. This result may have some theoretical implications which we are currently investigating.” In our view, the problem is not the finiteness of Z_S and Z_T , but whether these coefficients are all that is required to renormalise the SUSY current. In this respect the situation with supersymmetric Ward identities is not analogous to the one dealt in ref. [36], concerning chiral Ward identities and the normalisation Z_A of the axial current.

3. The Faddeev-Popov term is already $O(g_0^2)$ at tree-level, as remarked after (4.30). Therefore it contributes to the Ward identity to order g_0^2 and should appear in (4.30), where it is missing. In other words, the claim $Z_{FP}^{(0)} = 0$, made right after eq. (4.28) is unjustified.

4. Before (4.34) the definition $\Delta \equiv \mathcal{O}\nabla_\mu S_\mu(x) + \frac{\delta\mathcal{O}}{\delta\xi(x)}|_{\xi=0} + \mathcal{O}\frac{\delta S_{GF}}{\delta\xi(x)}|_{\xi=0} + \mathcal{O}B_i$ is not consistent with (4.31), (4.32), as it contains the terms $\mathcal{O}B_i$, whose second order contributions do not appear in (4.31), (4.32). The contributions from $\mathcal{O}B_i$ should not be included in (4.34).

5. The strategy for calculating the renormalisation constants without evaluating the one-loop contribution of the complicated term X_S requires knowledge of the list of mixing operators B_j appearing in (3.8). This implies an analysis of possible BRS-exact operators of dimension smaller than $\frac{11}{2}$. This analysis has not been carried out and it is not clear whether B_0 , B_1 , B_2 and B_3 form a complete set. For example, $B_4 = \frac{2}{g}(\partial_\rho A_\mu)\gamma_\mu\partial_\rho\lambda$, being independent of the other operators B_j , should have been considered at the general level of eq. (3.10), although its contribution vanishes at the specific kinematical regime of eq. (4.33).

6. The work presented in A. Feo’s paper is part of a joint project within the framework of the DESY-Münster-Roma collaboration. All involved researchers and students had agreed on joint publications of their results. The main contribution of A. Feo is the detailed implementation, programming and numerical evaluation of the various perturbative contributions. This considerable work, which we appreciate, is announced for publication in the sequel paper. The general procedure outlined in Feo’s present paper, however, grew out of discussions and internal written memoranda by the three of us. In Refs. [37,38,40] these sources are acknowledged as private communications and internal notes. None of us, however, has authorized their external use. Moreover, any definitive publication was postponed until the open questions (exposed in points 2. and 5. above) were resolved. Thus, so far we have limited ourselves to the joint presentation of partial results in past lattice workshops (see hep-lat/0110113, hep-lat/0011030).

We regret that last year A. Feo decided to cease collaborating with us. We express our total disapproval of this unauthorized publication of the collaboration’s internal documents

on unfinished work. We declare that we will refrain from any other public discussion or further comment on this matter, be it on the present preprint archive or elsewhere.